

## EVAPOTRANSPIRATION AND CROP COEFFICIENT FOR RADISH UNDER PROTECTED CULTIVATION

M. O. Pereira<sup>1</sup>, P. F. Silva<sup>2</sup>, R. M. Matos<sup>3</sup>, V. E. Borges<sup>4</sup>, T. Galvão Sobrinho<sup>5</sup>,  
J. Dantas Neto<sup>6</sup>

**ABSTRACT:** The objective of this study was to evaluate the real crop evapotranspiration and the crop coefficient of radish, “Crimson Gigante” cultivar, for different phenological stages under protected cultivation. The research was performed from March to April 2015 in pots arranged in greenhouse located at the Federal University of Campina Grande (UFCG). For plants irrigation it was used dripping irrigation system and the management was based on reference evapotranspiration ( $ET_0$ ). The water efficiency application of the irrigation system ( $E_a$ ), the gross irrigation depth (LB), the real crop evapotranspiration ( $ET_r$ ) and the crop coefficient ( $k_c$ ) were determined during the cycle cultivation of radish. The results showed  $E_a = 89,30\%$ , average daily LB of 4.31 mm, accumulated values of  $ET_0$  and  $ET_r$  of 175 and 150 mm. The maximum  $k_c$  (0.94) was obtained in the intermediate phenological stage (III) and the minimum  $k_c$  (0.58) occurred in the final stage (IV) with  $ET_r$  of 75.4 and 14.5 mm, respectively. The crop coefficients were similar to those in the literature, except in the last growth plant stage.

**KEYWORDS:** Crimson Gigante, greenhouse, water consumption.

## EVAPOTRANSPIRAÇÃO E COEFICIENTE DE CULTIVO PARA O RABANETE EM AMBIENTE PROTEGIDO

**RESUMO:** Objetivou-se no presente estudo, avaliar a evapotranspiração real e o coeficiente de cultivo do rabanete, cultivar Crimson Gigante, para as diferentes fases fenológicas da cultura em condições de ambiente protegido. A pesquisa foi conduzida de março a abril de 2015 em vasos dispostos em casa de vegetação na Universidade Federal de Campina Grande (UFCG). O sistema de irrigação foi por gotejamento e o manejo se baseou na evapotranspiração de referência ( $ET_0$ ). Foram determinadas a eficiência de aplicação de água do sistema de irrigação

<sup>1</sup> Doutoranda, UAEA/CTRN/UFCG. Campina Grande - Paraíba. email: marianapereira.agri@gmail.com;

<sup>2</sup> Pós-Doutoranda PPGRN/CTRN/UFCG. Campina Grande - Paraíba. Email: patrycyafs@yahoo.com.br;

<sup>3</sup> Doutorando UAEA/CTRN/UFCG. Campina Grande - Paraíba. Email: rigobertomoreira@gmail.com;

<sup>4</sup> Mestranda, UAEA/CTRN/UFCG. Campina Grande - Paraíba. email: kecis.borges@hotmail.com;

<sup>5</sup> Acadêmico, UAEA/CTRN/UFCG. Campina Grande - Paraíba. email: tgs\_galvao@hotmail.com;

<sup>6</sup> Prof. Dr, UAEA/CTRN/UFCG. Campina Grande - Paraíba. email: zedantas1955@gmail.com.

( $E_a$ ), a lâmina bruta de irrigação (LB), a evapotranspiração real da cultura ( $ET_r$ ) e o coeficiente de cultivo ( $k_c$ ) ao longo do ciclo do rabanete. Os resultados demonstraram  $E_a = 89,30\%$ , LB média  $4,31 \text{ mm dia}^{-1}$ , valores acumulados de  $ET_o$  e  $ET_r$  de 175 e 150 mm. O  $k_c$  máximo (0,94) foi obtido na fase fenológica intermediária (III) e o  $k_c$  mínimo (0,58) ocorreu na fase final (IV) com  $ET_r$  de, respectivamente, 75,40 e 14,50 mm. Os coeficientes de cultivo assemelharam-se aos existentes na literatura, exceto para a última fase de desenvolvimento da planta.

**PALAVRAS-CHAVE:** Crimson Gigante, casa de vegetação, consumo hídrico.

## INTRODUCTION

Radish (*Raphanus sativus* L.), an herbaceous plant belonging to the family Brassicaceae, is a small vegetable whose comestible part is the root. This plant is characterized as being very sensitive to variations in soil moisture and in a situation of minimal scarcity or excess water, presents physiological disturbances that interfere in its productivity and commercial root diameter, especially due to the cracks that causes in the tubercle (Filgueira, 2008).

Among the irrigation systems recommended for this crop, drip irrigation with high efficiency of water use in agriculture is evidenced, because it's applies the water irrigation depth only in the region of the root zone of the crops, maintaining the moisture of the soil close to the field capacity (Mantovani et al., 2012).

However, it is very important to know the water requirements of the radish crop so that an efficient management of the irrigation can be carried out and, in this way, the soil moisture conditions can be maintained so that the crop can satisfy its water needs during the different stages of development (Alves et al., 2017).

The parameter that reflects the water requirements by the plants is the evapotranspiration with substantial relevance for the irrigation, because it represents the amount of water necessary to be replaced by the irrigation (Camargo & Sentelhas, 1997). The evapotranspiration expresses the simultaneous occurrence of the processes of soil water evaporation and plant transpiration and it is a function of energy balance, atmospheric demand and soil water supply to plants (Adorian et al., 2015).

One of the procedures used to estimate crop water requirements involves the determination of reference evapotranspiration ( $ET_0$ ), which, through the use of appropriate crop coefficient ( $k_c$ ), allows the estimation of crop evapotranspiration ( $ET_c$ ) in the different crop development stages (Sediyama, 1987). In this way, the  $k_c$  is directly related to the phenological

and physiological stages of the crops and their respective water requirements, correlating them with the reference evapotranspiration ( $ET_0$ ). Crop evapotranspiration under real conditions of atmospheric factors and soil moisture characterizes the real crop evapotranspiration ( $ET_r$ ).

In this context, the objective of this study was to evaluate the real crop evapotranspiration and the crop coefficient of radish, “Crimson Gigante” cultivar, for different phenological stages under protected cultivation.

## MATERIAL AND METHODS

The research was performed in a greenhouse located in Campus I at Federal University of Campina Grande (UFCG), Campina Grande city (7° 13'50" S, 35° 52'52" W, 551 als), state of Paraíba, Brazil. The climate of the region is defined as As, tropical, with winter rains and dry summer, and average temperature of the hottest month above 28 °C (Coelho & Soncin, 1982).

The radish cultivar used was the “Crimson Gigante”, the seeds were produced in expanded polyethylene trays of 128 cells filled with commercial substrate. Transplanting occurred at 8 days after sowing. The crop was cultivated in 60 polyethylene pots with 12 L and with 1.0 m spacing between the rows and 0.50 m between plants. In each pot a 1 cm layer of gravel number 1 was placed, covered with geotextile blanket and 14 dm<sup>3</sup> of soil. To convert the pots into drainage lysimeters to perform water balance, a hole was made at base of each pot. The soil used was classified as medium texture according to Embrapa (2013) methodology.

In the research it was utilized a dripping irrigation system with a flow of 2.3 L h<sup>-1</sup> and the irrigation management was based on the Hargreaves & Samani (1982) model, according to methodology proposed by Medeiros et al. (2013) for reference evapotranspiration ( $ET_0$ ) determination (Equation 1):

$$ET_0 = 0,0023(T_{méd} + 17,8) * (T_{máx} - T_{mín})^{0,5}(R_a * 0,408) \quad (1)$$

where:

$ET_0$  – reference evapotranspiration, mm day<sup>-1</sup>

$T_{méd}$  – average temperature, °C

$T_{máx}$  – maximum temperature, °C

$T_{mín}$  – minimum temperature, °C

$R_a$  – extraterrestrial radiation, MJ m<sup>-1</sup> day<sup>-1</sup>

The air temperature and air relative humidity data during the experimental period from sowing until harvesting, equivalent to 37 days, were collected by sensors installed inside the greenhouse.

The Christiansen uniformity coefficient (CUC) of the irrigation system was 94 %, considering a potential efficiency of the irrigation system ( $E_{ap}$ ) of 95 % according to Equation 2 (Bernardo et al., 2008):

$$E_a = CUC * E_{ap} \quad (2)$$

where:

$E_a$  – water application efficiency of the irrigation system, %

$CUC$  – Christiansen uniformity coefficient, %

$E_{ap}$  – potential application efficiency, %

The gross irrigation depth was determined as a function of the crop evapotranspiration and the efficiency of the irrigation system through Equation 3 (Mantovani et al., 2012):

$$LB = \left( \frac{ET_c}{E_a} \right) * 100 \quad (3)$$

where:

$LB$  – gross irrigation depth, mm day<sup>-1</sup>

$ET_c$  – crop evapotranspiration, mm day<sup>-1</sup>

The real crop evapotranspiration ( $ET_r$ ) was quantified through the water balance (Silva et al., 2005), according to Equation 4:

$$\Delta ARM = P + I \pm R + AC - DP - ET_r \quad (4)$$

where:

$\Delta ARM$  – soil water storage variation, mm dia<sup>-1</sup>

$P$  – precipitation, mm dia<sup>-1</sup>

$I$  – irrigation, mm dia<sup>-1</sup>

$R$  – run off, mm dia<sup>-1</sup>

$AC$  – capillary water ascension, mm dia<sup>-1</sup>

$DP$  – deep drainage, mm dia<sup>-1</sup>

$ET_r$  – real crop evapotranspiration, mm dia<sup>-1</sup>

In the conditions of radish cultivation, the terms  $\Delta Arm$ ,  $P$ ,  $R$ ,  $AC$  and  $DP$  were considered null, because it is an experiment in pots, with frequent irrigation in greenhouse. Rearranging the terms of Equation 4, we have Equation 5:

$$ET_r = I \quad (5)$$

Thus, it was assumed that the  $ET_r$  value between two irrigation events was equal to the quantity of water applied to the soil in question, considering that the moisture was uniform throughout the soil profile. The crop coefficient ( $k_c$ ) was estimated according to Equation 6:

$$k_c = \frac{ET_r}{ET_0} \quad (6)$$

where,

$k_c$  – crop coefficient, dimensionless

The results of  $k_c$  were compared with the values found by Doorenbos & Pruitt (1977), Marouelli (2007), FAO-56 (Allen et al., 1998) for each phenological phase of the crop studied. The crop cycle according to Doorenbos & Pruitt (1977) comprises the initial stage in trays, phenological phase I; the second phase comprises the growth phase with duration characterized as phenological phase II; the third phase comprises the intermediate phase characterized as phenological phase III; and the final phase characterized as phenological phase IV.

## RESULTS AND DISCUSSION

The average air temperature inside the greenhouse was 29.6 °C with variation between 23.4 and 37.5 °C (Figure 1A). The average of air relative humidity was 59 % with variation between 41 and 79 % (Figure 1B).

The water application efficiency of the irrigation system was 89.30 % being classified as acceptable by Bernardo (2006) for dripping irrigation system. The average daily gross irrigation depth (LB) was 4.31 mm, corresponding to 91.31 % of reference evapotranspiration ( $ET_0$ ). Lacerda et al. (2017) verified that irrigation depth of 100 %  $ET_0$  and values close to this one have a better effect on the development and productivity of the crop, so the depth used in the present study was adequate.

The results of reference evapotranspiration ( $ET_0$ ) and real crop evapotranspiration ( $ET_r$ ) can be observed in Figure 2. Reference evapotranspiration showed average daily value of 4.72 mm and accumulated value close to 175 mm. While daily average value of real crop evapotranspiration was 4.05 mm with accumulated value next to 150 mm.  $ET_r$  accumulated value was 14 % less than accumulated of  $ET_0$ .

Oliveira Neto et al. (2011) in research with beet, also a tuberous root, cultivated under different dead coverages, observed that the maximum water consumption of the crop was 4.0 mm day<sup>-1</sup> and was close to the consumption of the radish of the present study.

The total water consumption of the crop, characterized by its evapotranspiration, for each phenological phase was 23.95 mm (phase I), 46.63 mm (phase II), 75.40 mm (phase III) and 14.50 mm phase IV). In daily average values, the ET<sub>r</sub> was 3.42 mm, 4.66 mm, 5.02 mm and 2.90 mm corresponding to the phases I, II, III e IV respectively. The duration of each phenological phase of the crop, in the order of the phases, was 7, 10, 15 and 5 days. The ET<sub>r</sub> increased with the development stages of the plant until the phenological stage III, from which it decreased in the final phase of the crop cycle (phenological phase IV) due to the senescence of the plants. This is because with the development of the crop, there is an increase of the water requirement by the plant to supply their physiological necessity, being phase III of filling of the tubercles the phase with the highest consumption.

Alves et al. (2017) in research with radish, “Crimson Gigante” cultivar, observed that the highest daily values of crop evapotranspiration were recorded in stage III (1.84 mm day<sup>-1</sup>). The stage IV presented the lowest water consumption because the culture was in the maturation stage and harvest of the fruits. These results are consistent with those obtained in the present work. However, ET<sub>r</sub> for the same phenological stage was 4.81 mm day<sup>-1</sup> and it was 2.6 times highest than the crop evapotranspiration found by Alves et al. (2017). This difference can be explained by the fact that climate is one of the main factors in determining the quantity of water evapotranspired by crops.

Radish crop coefficients ( $k_c$ ) for the different phenological phases are shown in figure 3. It was observed that the  $k_c$  of radish “Crimson Gigante” cultivar was higher in phenological phase III (0.94), followed by phenological phases II (0.84), I (0.73) and IV (0.58). Considering that the maximum  $k_c$  coincided with the phase of highest water consumption characterized by the highest ET<sub>r</sub> and that the same tendency occurred for the other development stages, it was verified the consistency of the results obtained for  $k_c$  values. When comparing the crop coefficients of the present study throughout the radish cycle with the values obtained by Doorenbos & Pruitt (1977), FAO-56 (Allen et al., 1998) and Marouelli (2007), it was verified that the coefficients obtained in this research for all phenological phases (Phase I = 0.70, Phase II = 0.80, Phase III = 0.90 and Phase IV = 0.55) were similar to the coefficients found by Marouelli (2007), followed by FAO-56 values with the exception of phase IV (phase I = 0.70, phase II = 0.80, phase III = 0.90 and phase IV = 0.85) (Figure 3). This variation of  $k_c$  values in relation to the others authors can be explained considering that the  $k_c$  was determined as a

function of  $ET_0$  that depends on the interaction between the various climatic elements (solar radiation, air temperature and air humidity) and that the  $ET_r$  was obtained under greenhouse conditions.

The coefficients obtained in the present study differed from the values found by Alves et al. (2017) for phenological phases I, II and IV (0.45, 0.55 and 0.65, respectively), but presented similarity in phase III (0.95). This fact is justified by the climatic differences of the places where the studies were conducted.

Silva et al. (2014) evaluated beet cultivar Early Wonder and verified maximum  $ET_r$  of  $2.37 \text{ mm day}^{-1}$  in the intermediate stage (phenological phase II) under salt stress conditions. For Itapuã 202's cultivar the highest  $ET_r$  was  $3.00 \text{ mm day}^{-1}$  at the same phase. These results differ from those obtained in the present study and can be justified because of salinity influences that can affect plant physiology and, in this way, reduce crop evapotranspiration. The authors also determined the crop coefficients for the different phenological phases of the beet and observed for phases I, II and III  $k_c$  values of approximately 0.3, 1.0 and 0.9 for Early Wonder and 0.20, 0.85 and 0.50 for the cultivar Itapuã 202. These differences can be attributed to local conditions, crop varieties and cultivation conditions.

## CONCLUSIONS

The water consumption and the crop coefficient of "Crimson Gigante" radish are maximum in the phenological phase III of development and formation of the tubercles and present minimum values in phase IV of maturation.

The crop coefficients are similar to those in the literature, except for the last stage of development of the plant (phenological phase IV).

The inside greenhouse climatic conditions affect the real crop evapotranspiration.

## REFERÊNCIAS BIBLIOGRÁFICAS

ADORIAN, G.C.; LORENCONI, R.; DOURADO NETO, D.; REICHARDT, K. Evapotranspiração potencial e coeficiente da cultura de dois genótipos de arroz de terras altas. Revista de Agricultura, V.90, n.2, p. 128-140, 2015.

ALLEN, R.G.; PEREIRA, L.S.; RAES, D.; SMITH, M. Crop evapotranspiration: guidelines for computing crop water requirements. Rome: FAO, 56, 1998, 297p.

ALVES, E.S.; LIMA, D.F.; BARRETO, J.A.S.; SANTOS, D.P.; SANTOS, M.A.L. Determinação do coeficiente de cultivo para a cultura do rabanete através de lisimetria de drenagem. *Irriga*, V.22, n.1, p. 194-203, 2017.

BERNARDO, S. Irrigação: total, suplementar, com déficit e de salvação. *Revista Irrigação & Tecnologia Moderna*, n.71/72, p. 30, 2006.

BERNARDO, S.; SOARES, A.; MANTOVANI, E.C. Manual de irrigação. 8. ed. Viçosa: UFV, Imprensa Universitária, 2008. 625p.

CAMARGO, Â.P.; SENTELHAS, P.C. Avaliação do desempenho de diferentes métodos de estimativa da evapotranspiração potencial no Estado de São Paulo, Brasil. *Revista Brasileira de Agrometeorologia*, V.5, n.1 p. 89-97, 1997.

COELHO, M.A.; SONCIN, N.B. Geografia do Brasil: São Paulo: Moderna, 1982. 368p.

SILVA, A.O.; FRANÇA, Ê.F.; KLAR, A.E.; CUNHA, A.R. Evapotranspiração e coeficiente de cultivo para a beterraba sob estresse salino em ambiente protegido. *Irriga*, V.19, n.3, p. 375, 2014.

DOORENBOS, J.; PRUITT, W.O. Guidelines for predicting crop water requirements. Rome: FAO, 1977. 179p. (Irrigation and Drainage Paper, 24).

FILGUEIRA, F.A.R. Novo manual de olericultura: agrotecnologia moderna na produção e comercialização de hortaliças. 3 ed. Viçosa: UFV, 2008. 422p.

HARGREAVES, G.H.; SAMANI, Z.A. Estimating potential evapotranspiration. *Journal of the Irrigation and Drainage Division*, V.108, n.3, p. 225-230, 1982.

MANTOVANI, E.C.; BERNARDO, S.; PALARETTI, L.F. Irrigação: princípios e métodos. 3 ed. atual e ampliada. Viçosa: UFV, 2012. 355p.

MAROUELLI, W. A. irrigação em campos de produção de sementes de hortaliças. Brasília: Empresa Brasileira de Pesquisa Agropecuária, 2007. 16p. (Circular Técnica, 52).

MEDEIROS, S.S.; REIS, C.F.; SANTOS JÚNIOR, J.A.; KLEIN, M.R.; RIBEIRO, M.D.; SZEKUT, F.D.; SANTOS, D.B. Manejo de irrigação utilizando o modelo de Hargreaves & Samani. INSA, Campina Grande. (Cartilha) 2013, 5p.

OLIVEIRA NETO, D.H.; CARVALHO, D.F.; SILVA, L.D.; GUERRA, J.G.M.; CEDDIA, M.B. Evapotranspiração e coeficientes de cultivo da beterraba orgânica sob cobertura morta de leguminosa e gramínea. *Horticultura Brasileira*, V.29, n.3, p. 330-334, 2011.




LACERDA, V.R.; GONÇALVES, B.G.; OLIVEIRA, F.G.; SOUSA, Y.B.; CASTRO, I.L. Características morfológicas e produtivas do rabanete sob diferentes lâminas de irrigação. *Revista Brasileira de Agricultura Irrigada-RBAI*, V.11, n.1, p. 1127-1134, 2017.

SEDIYAMA, G.C. Necessidade de água para os cultivos. In: ASSOCIAÇÃO BRASILEIRA DE EDUCAÇÃO AGRÍCOLA SUPERIOR, Brasília. Anais eletrônicos... Brasília, 1987. 143p.


SILVA, E.F.F.; CAMPECHE, L.F.S.M.C.; DUARTE, S.N.; FOLEGATTI, M.V. Evapotranspiração, coeficiente de cultivo e de salinidade para o pimentão cultivado em estufa. *Magistra*, V.17, n.2, p. 58-63, 2005.

EMBRAPA. Sistema brasileiro de classificação de solos. Rio de Janeiro: Empresa Brasileira Agropecuária, Centro Nacional de Pesquisa de Solos, 2013. 353p.


### Figure captions




**Figure 1A.** Air temperature over the experimental period.



**Figure 1B.** Air relative humidity over the experimental period.



**Figure 2.** Accumulated values of evapotranspiration (ET<sub>0</sub>) and real crop evapotranspiration (ET<sub>r</sub>) for “Crimson Gigante” cultivar.



**Figure 3.** Crop coefficient of “Crimson Gigante” cultivar for the different phenological phases.